arXiv:1703.04966v1 [gr-qc] 15 Mar 2017

General dynamical properties of cosmological models with nonminimal kinetic coupling

Jiro Matsumoto* and Sergey V. Sushkov†

Institute of Physics, Kazan Federal University,

Kremlevskaya Street 16a, Kazan 420008, Russia

Abstract

We consider cosmological dynamics in the theory of gravity with the scalar field possessing the nonminimal kinetic coupling to curvature given as $\eta G^{\mu\nu}\phi_{,\mu}\phi_{,\nu}$, where η is an arbitrary coupling parameter, and the scalar potential $V(\phi)$ which assumed to be as general as possible. With an appropriate dimensionless parameterization we represent the field equations as an autonomous dynamical system which contains ultimately only one arbitrary function $\chi(x) = 8\pi G|\eta|V(x/\sqrt{8\pi G})$. Then, assuming the rather general properties of $\chi(x)$, we analyze stationary points and their stability, as well as all possible asymptotical regimes of the dynamical system. It has been shown that for a broad class of $\chi(x)$ there exist attractors representing three accelerated regimes of the Universe evolution, including de Sitter expansion (or late-time inflation), the Little Rip scenario, and the Big Rip scenario. As the specific examples, we consider a power-law potential $V(\phi) = M^4(\phi/\phi_0)^\alpha$, Higgs-like potential $V(\phi) = \frac{\lambda}{4}(\phi^2 - \phi_0^2)^2$, and exponential potential $V(\phi) = M^4 e^{-\phi/\phi_0}$.

PACS numbers: 98.80.-k,95.36.+x,04.50.Kd

* E-mail address: jmatsumoto@kpfu.ru

† E-mail address: sergey_sushkov@mail.ru

I. INTRODUCTION

The current accelerated expansion of the Universe was clarified by the observations of Type Ia supernovae in 1990s [1, 2], and it is now also supported by the other observations: Cosmic Microwave Background Radiation (CMB) [3–5], Baryon Acoustic Oscillations (BAO) [6–10], and so on. Whereas, inflation hypothesis, which is the accelerated expansion of the Universe in early Universe, has been broadly investigated to explain the isotropic signals of CMB and the flatness problem. These accelerated expansion of the Universe cannot be described by the Einstein equations if we only take the usual matter into account. Therefore, introducing the cosmological constant or dynamical scalar field(s) into the equations or modifications of the Einstein gravity have been considered. In this paper, we treat a model which have a coupling between the kinetic term of the scalar field and the Einstein tensor [11–20]. This model can be regarded as a generalization of quintessence model [21], whereas, it can be also regarded as a special case of Horndeski's theory [22], which is known as a most general single scalar field model of dark energy that contains less than second derivatives in the field equations. The dynamical properties of the model has been investigated in the cases of a power-law potential [23], a Higgs-like potential [24], and so on. In this paper, we will consider the general potential $V(\phi)$, and analyze the field equations by regarding those as an autonomous system. The purposes of the research are not only to find some de Sitter-like solutions but also to clarify all the attractors that the model can describe. The contents of the paper are as follows; in Sec. II, we derive general equations in the Friedmann-Lemaitre-Robertson-Walker (FLRW) space-time, in Sec. III, the way to regard the field equations as an autonomous system is introduced, subsequently, the stability of the stationary points and the properties of the other attractors are investigated in detail, some concrete examples of the results derived in Sec. III are given in Sec. IV, concluding remarks are in Sec. V. The readers who are not so familiar with the dynamical analysis in this model are recommended to see the figures in Sec. IV before reading Sec. III because the calculations in Sec. III are so abstract. We use Natural units, $\hbar = c = k_{\rm B} = 1$, in the following.

II. ACTION AND FIELD EQUATIONS

Let us consider the theory of gravity with the action

$$S = \int d^4x\sqrt{-g}\left\{\frac{R}{8\pi G} - \left[g^{\mu\nu} + \eta G^{\mu\nu}\right]\phi_{,\mu}\phi_{,\nu} - 2V(\phi)\right\}, \tag{2.1}$$

where $V(\phi)$ is a scalar field potential, $g_{\mu\nu}$ is a metric, R is the scalar curvature, and $G_{\mu\nu}$ is the Einstein tensor. The coupling parameter η has the dimension of inverse mass-squared. Note that in the literature there is discussion, which is still open, about acceptable values of η. In Refs. [17, 18] it was assumed that $\eta < 0$ in order to prevent the appearance of ghosts in the model. However, analyzing scalar and tensor perturbations generated in the theories given by the action (2.1), Tsujikawa in Ref. [19] had derived more general conditions in order to avoid the appearance of scalar ghosts and Laplacian instabilities. Generally, their fulfillment depends on particular cosmological scenarios and is not directly determined by the sign of η. Moreover, in Ref. [20] it was demonstrated that the necessary condition that ensures the absence of instabilities is fulfilled for η both positive and negative. Since the detailed investigation of the instabilities and superluminality of the model with nonminimal kinetic coupling lies beyond the scope of the present work, hereinafter we will not impose any restrictions on values of η.

In the spatially-flat Friedmann-Robertson-Walker cosmological model the action (2.1) yields the following field equations [13]

$$3H^2 = 4\pi G\dot{\phi}^2\left(1 - 9\eta H^2\right) + 8\pi G V(\phi), \tag{2.2a}$$

$$2\dot{H} + 3H^2 = -4\pi G\dot{\phi}^2\left[1 + \eta\left(2\dot{H} + 3H^2 + 4H\ddot{\phi}\dot{\phi}^{-1}\right)\right] + 8\pi G V(\phi), \tag{2.2b}$$

$$(\ddot{\phi} + 3H\dot{\phi}) - 3\eta(H^2\ddot{\phi} + 2H\dot{H}\dot{\phi} + 3H^3\dot{\phi}) = -V_\phi, \tag{2.2c}$$

where a dot denotes derivatives with respect to time, $H(t) = \dot{a}(t)/a(t)$ is the Hubble parameter, $a(t)$ is the scale factor, $\phi(t)$ is a homogeneous scalar field, and $V_\phi = dV/d\phi$. Note that equations (2.2b) and (2.2c) are of second order, while (2.2a) is a first-order differential constraint for $a(t)$ and $\phi(t)$. The constraint (2.2a) can be rewritten as

$$\dot{\phi}^2 = \frac{3H^2 - 8\pi G V(\phi)}{4\pi G(1 - 9\eta H^2)}, \tag{2.3}$$

or equivalently as

$$H^2 = \frac{4\pi G\dot{\phi}^2 + 8\pi G V(\phi)}{3(1 + 12\pi G\eta\dot{\phi}^2)}. \tag{2.4}$$

Therefore, as long as the parameter η and the potential $V(\phi)$ are given, the above relations provide restrictions for the possible values of H and $\dot{\phi}$, since they have to give rise to non-negative $\dot{\phi}^2$ and H^2, respectively. Assuming the non-negativity of the potential, i.e. $V(\phi) \geq 0$, we can conclude from Eqs. (2.3) and (2.4) that in the theory with the positive η possible values of $\dot{\phi}$ are unbounded, while H takes restricted values. Vice versa, the negative η leads to bounded $\dot{\phi}$ and unbounded H.

Let us now resolve the equations (2.2b) and (2.2c) with respect to $\dot{H}$ and $\ddot{\phi}$, and then, using the relations (2.3) and (2.4), eliminate $\dot{\phi}$ and H from respective equations. As the result, we obtain

$$\dot{H} = \frac{-(1-3\eta H^2)(1-9\eta H^2)[3H^2 - 8\pi G V(\phi)] - 4\sqrt{\pi G}\eta H \sqrt{(1-9\eta H^2)[3H^2 - 8\pi G V(\phi)]}\, V_\phi}{1 - 9\eta H^2 + 54\eta^2 H^4 - 8\pi G \eta V(\phi)(1+9\eta H^2)}, \tag{2.5}$$

$$\begin{aligned}\ddot{\phi} =& \big[1 + 12\pi G\eta\dot{\phi}^2 + 96\pi^2 G^2\eta^2\dot{\phi}^4 + 8\pi G\eta V(\phi)(12\pi G\eta\dot{\phi}^2 - 1)\big]^{-1} \\ &\times \Big\{ -2\sqrt{3\pi G}\dot{\phi}[1 + 8\pi G\eta\dot{\phi}^2 - 8\pi G\eta V(\phi)]\sqrt{[\dot{\phi}^2 + 2V(\phi)](12\pi G\eta\dot{\phi}^2 + 1)} \\ &- (12\pi G\eta\dot{\phi}^2 + 1)(4\pi G\eta\dot{\phi}^2 + 1)V_\phi \Big\}. \end{aligned} \tag{2.6}$$

It is seen that the H-equation (2.5) in general contains ϕ-terms arising from the potential $V(\phi)$. At the same time, the ϕ-equation (2.6) does not contains H-terms, so that it represents a closed second-order differential equation for the unknown function $\phi(t)$. Such the form of Eq. (2.6) dictates us the following strategy of solving the system of field equations (2.2). First, analyzing the ϕ-equation (2.6), we will characterize in detail a time evolution of the scalar field $\phi(t)$. Then, using the constraint (2.4), we will be able to describe an evolution of the Hubble parameter $H(t)$.

III. SCALAR FIELD AS AN AUTONOMOUS DYNAMICAL SYSTEM

In this section we will focus on solving Eq. (2.6) which describes a time evolution of the scalar field.

Note that Eq. (2.6) has a normal form, i.e. resolved with respect to the second derivative. This allows us to use standard methods of the theory of dynamical systems. In practice, an analysis of Eq. (2.6) depends on particular values of the coupling parameter η, and hence further we will separately consider the cases $\eta > 0$, $\eta < 0$, and $\eta = 0$.

The case $\eta > 0$

Assume that η is positive. Now, let us introduce the set of dimensionless variables

$$x = \sqrt{8\pi G}\phi, \quad y = \sqrt{8\pi G\eta}\,\dot{\phi}, \quad \tau = \frac{t}{\sqrt{\eta}}, \quad \chi(x) = 8\pi G\eta V(x/\sqrt{8\pi G}). \tag{3.1}$$

By using dimensionless variables, we can rewrite Eq. (2.6) as the following autonomous dynamical system:

$$\frac{dx}{d\tau} = y, \tag{3.2}$$

$$\frac{dy}{d\tau} = \frac{-\sqrt{3}y\left[1+y^2-\chi(x)\right]\sqrt{\left[\frac{1}{2}y^2+\chi(x)\right]\left(\frac{3}{2}y^2+1\right)} - \left(\frac{3}{2}y^2+1\right)\left(\frac{1}{2}y^2+1\right)\chi'(x)}{1+\frac{3}{2}y^2+\frac{3}{2}y^4+\chi(x)\left(\frac{3}{2}y^2-1\right)}. \tag{3.3}$$

As seen in the system (3.2)-(3.3), the degrees of freedom of the model, which are the function $V(\phi)$ and the constant η in Eq. (2.6), are simply expressed as $\chi(x)$. Therefore, the mathematical properties of Eq. (3.2)-(3.3) only depend on $\chi(x)$, although the physical properties of the original equation (2.6) depend on both of $V(\phi)$ and η. Below we examine in detail basic features of the system (3.2)-(3.3).

1. Stationary points

Stationary points are those where $dx/d\tau = 0$ and $dy/d\tau = 0$. Hence, from Eq. (3.2) it follows that any stationary point of the system (3.2)-(3.3) has $y = 0$. Then, Eq. (3.3) yields

$$\left.\frac{dy}{d\tau}\right|_{y=0} = \frac{-\chi'(x)}{1-\chi(x)}. \tag{3.4}$$

The equality $(dy/d\tau)_{y=0} = 0$ is fulfilled if $\chi'(x) = 0$ and $\chi(x) \neq 1$.

The stability of stationary points with respect to small perturbations δx and δy can be investigated by fluctuating Eqs. (3.2)-(3.3) as

$$\left.\frac{d}{d\tau}\begin{pmatrix}\delta x\\ \delta y\end{pmatrix}\right|_{y=0,\chi'(x)=0} = \left.\begin{pmatrix}0 & 1\\ \frac{\partial(dy/d\tau)}{\partial x} & \frac{\partial(dy/d\tau)}{\partial y}\end{pmatrix}\right|_{y=0,\chi'(x)=0}\begin{pmatrix}\delta x\\ \delta y\end{pmatrix}. \tag{3.5}$$

A character of stationary points $(x, y) = (x_0, y_0)$ and a behavior of phase trajectories in their vicinity are determined by the eigenvalues of the matrix standing in the right-hand side of Eq. (3.5). In particular, (x_0, y_0) is stable if all real parts of eigenvalues are negative.

One can easily check that a 2×2 matrix $\{(0, a), (b, c)\}$ has, generally speaking, two eigenvalues $(c \pm \sqrt{c^2 + 4ab})/2$. Both of them have negative real parts if and only if $c < 0$ and $ab < 0$. Moreover, since a configuration of phase trajectories is determined by imaginary parts of eigenvalues, phase trajectories form spirals if $c^2 + 4ab < 0$, and straight lines if $c^2 + 4ab \geq 0$. Using these algebraic facts, we can conclude that, in our case, the stationary points (x_0, y_0) are stable if and only if

$$\left.\frac{\partial(dy/d\tau)}{\partial x}\right|_{(x,y)=(x_0,y_0)} < 0 \qquad \text{and} \qquad \left.\frac{\partial(dy/d\tau)}{\partial y}\right|_{(x,y)=(x_0,y_0)} < 0. \tag{3.6}$$

The concrete form of these conditions (3.6) are written down as

$$\frac{-\chi''(x_0)}{1-\chi(x_0)} < 0 \qquad \text{and} \qquad -\sqrt{3\chi(x_0)} < 0, \tag{3.7}$$

respectively. The second equation of (3.7) is always satisfied if $\chi(x_0) > 0$, whereas, the first equation of (3.7) implies that the stability of the stationary point depends on the value of η when the value of the potential does not vanish. Therefore, there are the case that η make the potential minimum unstable and the case that η make the potential maximum stable if $V(\phi_0 = x_0/\sqrt{8\pi G}) \neq 0$.

While, in the case of $V(\phi_0) = 0$, we need to carefully deal with the stability conditions, because the second equation of (3.7) vanishes. Zero real parts of the eigenvalues mean that the stability cannot be understood only from the linear perturbation terms around the stationary point. The next leading order terms, i.e. second order terms should be taken into account. Expanding Eqs. (3.2) and (3.3) around $(x, y) = (x_0, 0)$ by applying $x \to x_0 + \delta x$ and $y \to 0 + \delta y$ yields

$$\frac{d\delta x}{d\tau} = \delta y, \tag{3.8}$$

$$\frac{d\delta y}{d\tau} = -\chi''(x_0)\delta x - \frac{1}{2}\chi'''(x_0)\delta x^2 - \sqrt{\frac{3}{2}\left[\delta y^2 + \chi''(x_0)\delta x^2\right]}\delta y + O(\delta^3), \tag{3.9}$$

where $\delta \sim \delta x \sim \delta y$. The δy dependence of $dy/d\tau$ is first appeared as the $O(\delta^2)$ terms. We can find Eqs. (3.8) and (3.9) do not depend on the value of η if we use the original variables. Therefore, these equations are equivalent to those in case of $\eta = 0$, i.e. the theory without kinetic coupling. The leading term of the right-hand side of the Eq. (3.9) with respect to δx is $-\chi''(x_0)\delta x$, and that with respect to δy is $-2\sqrt{3(\chi''(x_0)\delta x^2 + \delta y^2)}\delta y$. By taking into account these terms, one can obtain the following approximated solution [25]:

$$\delta x = \sqrt{\frac{8}{3\chi''(x_0)}}\frac{1}{\tau}\sin\left(\sqrt{\chi''(x_0)}\tau\right) = \sqrt{\frac{8}{3V''(\phi_0)}}\frac{1}{t}\sin\left(\sqrt{V''(\phi_0)}t\right). \tag{3.10}$$

Therefore, the stationary point which satisfies $\chi(x_0) = \chi'(x_0) = 0$ and $\chi''(x_0) > 0$ is stable. If $\chi''(x_0) < 0$, the point $(x, y) = (x_0, 0)$ is unstable as seen from the first equation of (3.7). Further calculations are required for the case $\chi(x_0) = \chi'(x_0) = \chi''(x_0) = 0$, however, we will not treat this case in this paper.

2. Critical curves

Generally speaking, there are many kinds of attractors in the phase diagram, e.g. stable stationary points, stable limit cycles, and the points located on infinity. In the following, we will consider "critical curves", $dy/d\tau = 0$ and $dy/d\tau = \pm\infty$, to investigate the behavior of the trajectories in the region $|x| \gg 1$ or $|y| \gg 1$. The autonomous system (3.2)-(3.3) show that $dx/d\tau$ is always positive in the first and second quadrant in the x-y phase diagram and $dx/d\tau < 0$ is satisfied in the third and fourth quadrants. Therefore, the attractors located on infinity only exist in the first and third quadrants. Then, the attractors are found by taking the sign of $dy/d\tau$ into account. The Poincaré compactification may be useful to investigate that region, however, we will not apply it for the system (3.2)-(3.3). Because not only the stability of the stationary points but also the varieties of the asymptotes are worthy to investigate.

The sufficient conditions for $dy/d\tau = \pm\infty$ are $\chi(x) \to \pm\infty$, $\chi'(x) \to \pm\infty$, or

$$1 + \frac{3}{2}y^2 + \frac{3}{2}y^4 + \chi(x)\left(\frac{3}{2}y^2 - 1\right) = 0, \tag{3.11}$$

if x and y are finite. The condition $|y| < \sqrt{2/3}$ is necessary to satisfy Eq. (3.11) when $\chi = 8\pi G\eta V$ is positive. If $\chi(x)$ continuously goes to infinity in the limit $x \to \pm\infty$, then the curves (3.11) asymptotically approach $y = \pm\sqrt{2/3}$. Whereas, solving Eq. (3.11) with respect to $\chi(x)$ gives a constraint $1 \leq \chi(x) < +\infty$. This result says that if η is small enough to satisfy $8\pi G\eta V(\phi) < 1$ for all ϕ, then there are no critical curves satisfying $dy/d\tau = \pm\infty$ except for the case $\chi(x) \to \pm\infty$ or $\chi'(x) \to \pm\infty$, conversely, if η is large enough to satisfy $8\pi G\eta V(\phi) \geq 1$ for some ϕ, then the critical curves appear. The curve (3.11) divide the whole phase diagram into several isolated areas, so it is essential to investigate Eq. (3.11) in this model. The trajectories in an isolated area cannot cross over the curve $dy/d\tau = \pm\infty$ without singularities or indefinableness of the physical quantities.

Region	Range of y	$\chi(x \to \pm\infty)$	Sign of $y\chi'$	Existence conditions	Nullclines
$y^2 \gg \chi$	$\lvert y\rvert \gg 1$	infinite	$-$	$\lvert\chi'\rvert \gg \chi^{1/2}$	$y = -\chi'/2$
$y^2 \sim \chi$	$\lvert y\rvert \gg 1$	infinite	$\pm$	$\lvert\chi'\rvert \sim \chi^{1/2}$	Eq. (3.18)
		infinite	$\pm$	$\lvert\chi'\rvert \ll \chi^{1/2}$	$y = \pm\sqrt{\chi}$
	$y = O(1)$	finite	0	$\chi > 1$	$y = \pm\sqrt{\chi - 1}$
$y^2 \ll \chi$	$\lvert y\rvert \gg 1$	infinite	$+$	$\chi^{1/2} \ll \lvert\chi'\rvert \ll \chi^{3/2}$	$y^3/\lvert y\rvert = 2\sqrt{2\chi^3}/\chi'$
	$y = O(1)$	infinite	$+$	$\lvert\chi'\rvert \sim \chi^{3/2}$	Eq. (3.16)
	$\lvert y\rvert \ll 1$	infinite	$+$	$\lvert\chi'\rvert \ll \chi^{3/2}$	$y = \chi'/\sqrt{3\chi^3}$
	$\lvert y\rvert \ll 1$	finite	0	$\chi \neq 1$	$y = -\chi'/[\sqrt{3\chi}(1-\chi)]$

Table I: Nullclines in the region $|x| \gg 1$ or $|y| \gg 1$. The signature of $y\chi'$ implies that the existence regions of the nullclines are constrained. For example, the continuous functions $\chi(x) > 0$ and $\chi'(x)$ cannot satisfy the conditions $\chi(+\infty) \gg 1$ and $y\chi'(+\infty) > 0$ in the fourth quadrant. Therefore, the nullclines of $y\chi' > 0$ only exist in the first and third quadrants.

The curve $dy/d\tau = 0$, which is called the nullcline, is expressed as follows:

$$-\sqrt{3}y\left[1 + y^2 - \chi(x)\right]\sqrt{\left[\tfrac{1}{2}y^2 + \chi(x)\right]\left(\tfrac{3}{2}y^2 + 1\right)} - \left(\frac{3}{2}y^2 + 1\right)\left(\frac{1}{2}y^2 + 1\right)\chi'(x) = 0. \quad (3.12)$$

In general, resolving Eq. (3.12) with respect to y is impossible. However, we can consider the asymptote of the nullcline (3.12) in the limit $x \to \pm\infty$ or $y \to \pm\infty$. In the following, $\chi(x)$ is treated as a nonnegative function.

First, the case $|x| \gg 1$ is investigated. There are three types of continuous function $\chi(x)$ in the limit $x \to \pm\infty$; that has a singular behavior, converging, or oscillating. Here, we will consider the first case and the second case. If $\chi(x)$ converges in the limit $x \to \pm\infty$, then $\chi'(x \to \pm\infty) = 0$ is satisfied. At this time, Eq. (3.12) yields $y = 0$ or $1 + y^2 = \chi$. The asymptote approaching the point $(x, y) = (\pm\infty, 0)$ is expressed as

$$y = -\frac{\chi'}{\sqrt{3\chi}(1-\chi)}, \quad (3.13)$$

whereas, $1 + y^2 = \chi$ means y converges into the finite value

$$y = \pm\sqrt{\chi(x \to \pm\infty) - 1} \quad (3.14)$$

if $\chi(x \to \pm\infty) > 1$. If $\chi(x)$ goes to infinity in the limit $x \to \pm\infty$, we can consider several asymptotes of $dy/d\tau = 0$. First, let us consider the case $1 \ll y^2 \ll \chi(x)$ as a case analysis,

then we have an asymptote of the nullcline,

$$y^2 = 2\sqrt{2}\frac{\chi^{3/2}(x)}{\chi'(x)}, \tag{3.15}$$

if $y\chi' > 0$ and $|\chi'| > \sqrt{\chi}$. Here, the condition $y\chi' > 0$ is imposed because the terms in Eq. (3.12) should be canceled, and the condition $|\chi'| > \sqrt{\chi}$ comes from the assumption $1 \ll y^2 \ll \chi(x)$.

When y is finite but $\chi \gg 1$ and $y\chi' > 0$, the asymptote

$$\sqrt{3\chi^3}y - \sqrt{\frac{3}{2}y^2 + 1}\left(\frac{1}{2}y^2 + 1\right)\chi' = 0, \tag{3.16}$$

is given if $\chi(x)^3/\chi'^2(x) \gtrsim 1$. If $\chi(x)^3/\chi'^2(x) \ll 1$ there is no real y satisfying $dy/d\tau = 0$ and $\chi \gg 1$. In particular, if the condition $\chi(x)^3/\chi'^2(x) \gg 1$ holds, then we have

$$y^2 = \frac{\chi'^2(x)}{3\chi^3(x)}. \tag{3.17}$$

In the case $1 \ll y^2 \sim \chi(x)$, the nullcline exists only if $|\chi'(x \to \pm\infty)| \lesssim \sqrt{\chi(x \to \pm\infty)}$, and is expressed as

$$(y^2 - \chi(x))\sqrt{\frac{1}{2}y^2\left(\frac{1}{2}y^2 + \chi(x)\right)} + \frac{1}{4}y^3\chi'(x) = 0. \tag{3.18}$$

In particular, the nullcline (3.18) is simplified for the case $|\chi'| \ll \sqrt{\chi}$ as

$$y = \pm\sqrt{\chi(x)}. \tag{3.19}$$

In the case $\chi(x) \ll y^2$ and $1 \ll y^2$, we have a nullcline

$$y = -\frac{1}{2}\chi'(x), \tag{3.20}$$

only if $y\chi' < 0$. The expression of the nullcline (3.20) implies that $\chi'(x)$ must satisfy $|\chi'| \gg 1$. Therefore, there is no nullcline which satisfy $1 \ll y^2$ and $\chi(x)$ is finite if $\chi(x)$ is a continuous function, moreover, there is no nullcline in the region $1 \ll y^2$ and x is finite if $\chi'(x)$ is a continuous function. All of the nullclines derived in this paragraph are expressed in Table I.

3. Behavior of trajectories in regions separated by the critical curves

First, we can find there are no attractors in the region $\chi^{1/2}, |\chi'| \ll |y|$ and $1 \ll |y|$, because the signature of $dy/d\tau$ is opposite to that of y. It can be also seen that the attractors in the

Type of the function $\chi(x)$		Asymptotes	Existence conditions	Hubble rate
$\chi(\pm\infty)$ is finite		$y=-\chi'/[\sqrt{3\chi}(1-\chi)]$	$\chi(\pm\infty)\neq 1$, $\chi'/(1-\chi)<0$	$H=\sqrt{\frac{8\pi GV}{3}}$
		$y=\pm\sqrt{\chi-1}$	$\chi(\pm\infty)>1$	$H=\frac{1}{\sqrt{3\eta}}$
$\chi(\pm\infty)$ is infinite	$\lvert\chi'\rvert\ll\chi^{1/2}$	$y=\frac{\chi'}{\sqrt{3\chi^3}}$		$H=\sqrt{\frac{8\pi GV}{3}}$
		$y=\pm\sqrt{\chi}$		$H=\frac{1}{\sqrt{3\eta}}$
	$\lvert\chi'\rvert\sim\chi^{1/2}$	$y=\frac{\chi'}{\sqrt{3\chi^3}}$		$H=\sqrt{\frac{8\pi GV}{3}}$
		$y=\alpha\sqrt{\chi}$, $\alpha\neq 0,\pm 1$		$H=\sqrt{\frac{2+\alpha^2}{9\alpha^2\eta}}$
	$\chi^{1/2}\ll\lvert\chi'\rvert\ll\chi^{3/2}$	$y=\frac{\chi'}{\sqrt{3\chi^3}}$		$H=\sqrt{\frac{8\pi GV}{3}}$
		$y^2=\alpha\frac{\sqrt{\chi^3}}{\chi'}$, $\alpha\neq 0,\pm 2\sqrt{2}$		$H=\sqrt{\frac{2V_\phi}{9\alpha(\kappa V)^{1/2}}}$
	$\lvert\chi'\rvert\sim\chi^{3/2}$	$y=c_1, 0<\lvert c_1\rvert<\sqrt{\frac{2}{3}}$	$0<\lvert\frac{\chi'}{\chi^{3/2}}\rvert<\frac{3}{4}$	$H=\sqrt{\frac{16\pi GV}{6+9c_1^2}}$
		$y=c_2, \sqrt{\frac{2}{3}}<\lvert c_2\rvert$	$0<\lvert\frac{\chi'}{\chi^{3/2}}\rvert<\frac{3}{4}$	$H=\sqrt{\frac{16\pi GV}{6+9c_2^2}}$
		$y=\pm\sqrt{\frac{2}{3}}$	$\frac{3}{4}<\lvert\frac{\chi'}{\chi^{3/2}}\rvert$	$H=\sqrt{\frac{4\pi GV}{3}}$
	$\chi^{3/2}\ll\lvert\chi'\rvert$	$y=\pm\sqrt{\frac{2}{3}}$		$H=\sqrt{\frac{4\pi GV}{3}}$

Table II: Asymptotes of the trajectories in the region $|x|\gg 1$.

limit $x\to\pm\infty$ or $y\to\pm\infty$ are only located in the region $x\to+\infty$ and $y>0$ and the region $x\to-\infty$ and $y<0$ by taking into account the signature of $dx/d\tau$. In the following, we will investigate the behaviors of the trajectories in the regions $x\gg+1$, $y>0$ and $x\ll-1$, $y<0$ by using case analysis with respect to the form of the function $\chi(x)$ in the limit $x\to\pm\infty$.

a. The case $\chi(\pm\infty)$ is finite When the function $\chi(x)$ is finite in the limit $x\to\pm\infty$, there are only nullclines $y_{n\pm}=\pm\sqrt{\chi-1}$ and $y=-\chi'/[\sqrt{3\chi}(1-\chi)]$ in the region $|x|\gg 1$ or $|y|\gg 1$ if we assume a continuity of $\chi(x)$ and $\chi'(x)$ as shown in the last subsection. The critical curves (3.11) are written down as

$$y_{s\pm}=\pm\sqrt{-\frac{1+\chi}{2}+\frac{1}{2}\sqrt{(1+\chi)^2-\frac{8}{3}(1-\chi)}} \tag{3.21}$$

in that region. The existence conditions for the nullclines $y_{n\pm}=\pm\sqrt{\chi-1}$ and it for the critical curves (3.21) are same, i.e. $\chi>1$. The absolute value of $y_{n\pm}$ is always larger than that of $y_{s\pm}$, so the signature of $dy/d\tau$ in the case $y>0$ is determined as follows; $dy/d\tau$ is negative above $y_{n+}=\sqrt{\chi-1}$, it becomes positive in the region $y_{s+}<y<y_{n+}$, and it

becomes negative if $-\chi'/[\sqrt{3\chi}(1-\chi)] < y < y_{s+}$. By taking into account the signatures of $dy/d\tau$ and $dx/d\tau$, we can find the nullcline $y_{n+} = +\sqrt{\chi - 1}$ is an attractor in the limit $x \to +\infty$, and $y_{n-} = -\sqrt{\chi - 1}$ is an attractor in the limit $x \to -\infty$. The stability of the curve $y = -\chi'/[\sqrt{3\chi}(1-\chi)]$ is determined by the sign of $\chi'/(1-\chi)$ because it determine the sign of y in the region $|x| \gg 1$. The condition that the curve should be in the first or third quadrant yields the stability condition $\chi'/(1-\chi) < 0$. The value of the Hubble rate function on $y_{n\pm} = \pm\sqrt{\chi - 1}$ and $y = 0$ are given as $H = 1/\sqrt{3\eta}$ and $H = \sqrt{8\pi GV/3}$, respectively.

b. The case $\chi(\pm\infty)$ is infinite and $|\chi'(\pm\infty)| \ll \chi^{1/2}(\pm\infty)$. If the functions $\chi(x)$ and $\chi'(x)$ are continuous functions, then $\chi'(+\infty) > 0$ ($\chi'(-\infty) < 0$) holds when $\chi(+\infty) \to +\infty$ ($\chi(-\infty) \to +\infty$) is satisfied. Therefore, we will consider the cases $\chi'(+\infty) > 0$ and $\chi'(-\infty) < 0$ in the following. If $\chi(\pm\infty)$ is infinite and $|\chi'(\pm\infty)|$ is much less than $\chi^{1/2}(\pm\infty)$, there are two kinds of nullclines in the limit $x \to \pm\infty$. The nullcline $y = \pm\sqrt{\chi(x)}$ is located in the region $y^2 \sim \chi(x)$ and the nullcline $y = \chi'(x)/\sqrt{3\chi^3(x)}$ is located in the region $y^2 \ll \chi(x)$. On the other hand, the critical curves (3.11) are approximately written down as $y = \pm\sqrt{2/3}$ in the region $|x| \gg 1$. In the right half of the x-y phase diagram, these critical curves are located in order from the top as $y = \sqrt{\chi(x)}$, $y = \sqrt{2/3}$, $y = \chi'(x)/\sqrt{3\chi^3(x)}$, $y = -\sqrt{2/3}$, and $y = -\sqrt{\chi(x)}$. The signature of $dy/d\tau$ is $-$ in the range $y > \sqrt{\chi(x)}$, is $+$ in $\sqrt{2/3} < y < \sqrt{\chi(x)}$, is $-$ in $\chi'(x)/\sqrt{3\chi^3(x)} < y < \sqrt{2/3}$, and so on.

By taking that $\chi'(x)/\sqrt{3\chi^3(x)} > 0$ and $\chi'(x)/\sqrt{3\chi^3(x)} \simeq const.$ hold in the region $|x| \gg 1$ into account, it is seen that $y = \chi'(x)/\sqrt{3\chi^3(x)}$ is an asymptote of the trajectories. The Hubble rate function on this asymptote is given by $H = \sqrt{8\pi GV/3}$. Whereas, the nullcline $y = \sqrt{\chi}$ is also an asymptote of the trajectories in the first quadrant and in the third quadrant as seen in the following. The inclination of y with respect to x of the nullcline $y = \pm\sqrt{\chi}$ is given as $dy/dx = \chi'/(2\sqrt{\chi}) \to 0$, whereas, the inclination of the trajectories on the curve $y = \alpha\sqrt{\chi}$ is $(dy/d\tau)/(dx/d\tau) = 2(1-\alpha^2)\sqrt{\alpha^2(\alpha^2+2)}/(3\alpha^5)$. Therefore, if the trajectories are once on the nullcline $y = \pm\sqrt{\chi}$, the trajectories continue being on it. At the same time, the condition $|(dy/d\tau)/(dx/d\tau)| \gtrsim 1$ is satisfied in the adjoining regions $1 \ll |y| \ll \sqrt{\chi}$ and $\sqrt{\chi} \ll |y|$, so the trajectories that are far from $y = \pm\sqrt{\chi}$ will approach the nullcline $y = \pm\sqrt{\chi}$. The Hubble rate function on this nullcline is given by $H = 1/\sqrt{3\eta}$.

c. The case $\chi(\pm\infty)$ is infinite and $|\chi'(\pm\infty)| \sim \chi^{1/2}(\pm\infty)$. The behavior of the trajectories in the case that $\chi(\pm\infty)$ is infinite and $|\chi'(\pm\infty)| \sim \chi^{1/2}(\pm\infty)$ is almost same as that we considered in the last paragraph. However, we should take care of the behaviors of

the trajectories in the region $y^2 \sim \chi(x)$. In general, the nullcline in the region $y^2 \sim \chi(x)$ is obtained by solving Eq. (3.18). If we only evaluate the leading part of the terms, we can treat χ' as $\chi' \propto \chi^{1/2}$ because $|\chi'(\pm\infty)| \sim \chi^{1/2}(\pm\infty)$. Then, we have $y = \alpha\chi^{1/2}$ as a solution of Eq. (3.18), where $\alpha \neq 0, 1$ is a real number. There is only one value of α that can satisfy Eq. (3.18) if we treat χ' as $\chi' = \beta\chi^{1/2}$, $\beta \in \mathbb{R}_{\neq 0}$. On the nullclines, $(dy/d\tau)/(dx/d\tau) = 0$ is satisfied because $dy/d\tau = 0$, whereas, the derivative of the nullcline $y = \alpha\chi^{1/2}$ yields $dy/dx \neq 0$. Therefore, the nullcline $y = \alpha\chi^{1/2}$ is not an asymptote of the trajectories. At the same time, if we consider the curve $y = f(x) = \gamma\chi^{1/2}$, where $\gamma \neq 0, 1, \alpha$ is a real number, we have $(dy/d\tau)/(dx/d\tau)|_{y=f(x)} = O(1)$ and $df(x)/dx = O(1)$. Therefore, we can find the asymptotes by solving the equation $(dy/d\tau)/(dx/d\tau)|_{y=f(x)} = df(x)/dx$ with respect to γ. There is only one value of γ that can satisfy this equality. The Hubble rate function on the asymptote is expressed as $H = \sqrt{(2+\gamma^2)/(9\gamma^2\eta)}$.

d. The case $\chi(\pm\infty)$ is infinite and $\chi^{1/2}(\pm\infty) \ll |\chi'(\pm\infty)| \ll \chi^{3/2}(\pm\infty)$. If $\chi(\pm\infty)$ is infinite and $\chi^{1/2}(\pm\infty) \ll |\chi'(\pm\infty)| \ll \chi^{3/2}(\pm\infty)$, there are three kinds of nullclines; $y = \chi'/\sqrt{3\chi^3}$, $y^3/|y| = 2\sqrt{2\chi^3}/\chi'$, and $y = -\chi'/2$. The behavior of the trajectories around the first one has been already considered in the other cases, and it is not changed in this case. The second one is located in the region $y^2 \ll \chi$ if $y\chi' > 0$, while, the third one is located in the region $y^2 \gg \chi$ if $y\chi' < 0$. In the region $2/3 < y^2 \ll 2\sqrt{2\chi^3}/|\chi'|$, the condition $(dy/d\tau)/(dx/d\tau) \gg 1$ is always satisfied, moreover, this inclination is more than that of the nullclines $y^3/|y| = 2\sqrt{2\chi^3}/\chi'$, so the trajectories in this region approach the nullclines $y^3/|y| = 2\sqrt{2\chi^3}/\chi'$.

In the region $y^2 \sim \chi$, the direction of the trajectories depends on the signature of $y\chi'$ as $(dy/d\tau)/(dx/d\tau) \sim -\chi'/y$. If $y\chi' > 0$, there are only nullclines $y^3/|y| = 2\sqrt{2\chi^3}/\chi'$ for $y^2 > 2/3$ and the inclination of the trajectories is negative in the region $y^2 \sim \chi$, so the curves $y^3/|y| = 2\sqrt{2\chi^3}/\chi'$ look stable when viewed from a distance. However, they are not attractors because the inclement of them do not vanish although $dy/d\tau = 0$ is satisfied on them. The attractors are, in fact, given by $y^2 = c_1\sqrt{\chi^3}/\chi'$, where $c_1 \neq \pm 2\sqrt{2}$ is a real number. The value of c_1 is exactly determined if the index i and the coefficient c of $\chi'(\pm\infty) = c\chi^i(\pm\infty)$ is given. The difference between the nullclines and the attractor is only the coefficient of the terms. On the other hand, if $y\chi' < 0$, there are only nullclines $y = -\chi'/2$ for $y^2 > 2/3$, however, they are only located in the second quadrant and in the fourth quadrant if χ is positive. Therefore, the nullclines $y = -\chi'/2$ have nothing to do with

the attractors in $x \to \pm\infty$.

e. The case $\chi(\pm\infty)$ is infinite and $|\chi'(\pm\infty)| \sim \chi^{3/2}(\pm\infty)$. If $\chi(\pm\infty)$ is infinite and $|\chi'(\pm\infty)| \sim \chi^{3/2}(\pm\infty)$, there are two kinds of nullclines; $y = -\chi'/2$ and Eq. (3.16). The nullclines $y = -\chi'/2$ have nothing to do with the attractors in the region $x \to \pm\infty$, whereas, Eq. (3.16) gives the characteristic configuration of the attractors. By assuming the function $\chi'(x)$ as $\chi'(x) = \alpha\chi^{3/2}(x)$ in the limit $x \to \pm\infty$, Eq. (3.16) gives

$$\alpha = \frac{\sqrt{3}y}{\sqrt{\frac{3}{2}y^2 + 1}\left(\frac{1}{2}y^2 + 1\right)}. \tag{3.22}$$

If $0 < |\alpha| < 3/4$, then there are two solutions in Eq. (3.22) and the range of the solutions are given by $0 < |y| < \sqrt{2/3}$ and $|y| > \sqrt{2/3}$, respectively. If $\alpha = \pm 3/4$, then there is one solution $y = \pm\sqrt{2/3}$, else if $|\alpha| > 3/4$, then there is no real solutions. Taking the existence of the critical curves $y = \pm\sqrt{2/3}$ into account gives the existence condition for the nullclines $y = const.$ as $0 < |\alpha| < 3/4$. The nullclines $y = const.$ are always stable for y direction, however, are only stable in the case $\chi'(+\infty) > 0$ or $\chi'(-\infty) < 0$ for x direction, therefore, they are the attractors if they are in the first quadrant or in the third quadrant. The case $|\alpha| > 3/4$ is same as the following case; there are no nullclines in the first and the third quadrant and the critical lines $y = \pm\sqrt{2/3}$ become stable.

f. The case $\chi(\pm\infty)$ is infinite and $\chi^{3/2}(\pm\infty) \ll |\chi'(\pm\infty)|$. There are only the nullclines $y = \chi'/2$ in the limit $x \to \pm\infty$ in the case $\chi(\pm\infty)$ is infinite and $\chi^{3/2}(\pm\infty) \ll |\chi'(\pm\infty)|$. Therefore, the critical curves that have a relation with the attractors are only $y = \pm\sqrt{2/3}$. In fact, $y = \sqrt{2/3}$ is the attractor in the limit $x \to +\infty$ and $y = -\sqrt{2/3}$ is the attractor in the limit $x \to -\infty$, although $dy/d\tau$ becomes singular on the curves $y = \pm\sqrt{2/3}$.

The asymptotes in the region $|x| \gg 1$ of the case $\eta > 0$ are summarized in Table II.

The case $\eta < 0$

In the case when η is negative, we need to redefine the set of dimensionless variables as follows

$$x = \sqrt{8\pi G}\phi, \quad y = \sqrt{8\pi G|\eta|}\,\dot{\phi}, \quad \tau = \frac{t}{\sqrt{|\eta|}}, \quad \chi(x) = 8\pi G|\eta|V(x/\sqrt{8\pi G}). \tag{3.23}$$

Now, we can rewrite Eq. (2.6) as the following autonomous dynamical system:

$$\frac{dx}{d\tau} = y, \tag{3.24}$$

Type of the function $\chi(x)$		(x,y) of the attractors	Stability conditions	Hubble rate
$\chi(\pm\infty)$ is finite		$(x_0,0)$, s.t. $\chi'(x_0)=0$	$\chi''(x_0)>0$	$H=\sqrt{\frac{8\pi GV}{3}}$
		$(\pm\infty,0)$	$\chi'(\pm\infty)=\mp 0$	$H=\sqrt{\frac{8\pi GV}{3}}$
$\chi(\pm\infty)$ is infinite	$\lvert\chi'\rvert \ll \chi^{3/2}$	$(x_0,0)$, s.t. $\chi'(x_0)=0$	$\chi''(x_0)>0$	$H=\sqrt{\frac{8\pi GV}{3}}$
	$\lvert\chi'\rvert \sim \chi^{3/2}$	$(x_0,0)$, s.t. $\chi'(x_0)=0$	$\chi''(x_0)>0$	$H=\sqrt{\frac{8\pi GV}{3}}$
		$(-\infty,y_0)$, $-\sqrt{\frac{2}{3}}<y_0<0$	$\chi(-\infty)\to+\infty$	$H=\sqrt{\frac{16\pi GV}{3(2-3y_0^2)}}$
	$\chi^{3/2} \ll \lvert\chi'\rvert$	$(x_0,0)$, s.t. $\chi'(x_0)=0$	$\chi''(x_0)>0$	$H=\sqrt{\frac{8\pi GV}{3}}$
		$(-\infty,-\sqrt{\frac{2}{3}})$	$\chi(-\infty)\to+\infty$	$H=+\infty$

Table III: Attractors in the case $\eta<0$.

$$\frac{dy}{d\tau}=\frac{-\sqrt{3}y\left[1-y^2+\chi(x)\right]\sqrt{\left[\frac{1}{2}y^2+\chi(x)\right]\left(1-\frac{3}{2}y^2\right)}-\left(1-\frac{3}{2}y^2\right)\left(1-\frac{1}{2}y^2\right)\chi'(x)}{1-\frac{3}{2}y^2+\frac{3}{2}y^4+\chi(x)\left(\frac{3}{2}y^2+1\right)}. \quad (3.25)$$

Here there is some difference from Eqs. (3.2) and (3.3) in the signs of the terms. The noteworthy points are that the definable range of y is constrained and the denominator of Eq. (3.25) cannot vanish. Therefore, the x-y phase diagram is quite different from that of $\eta>0$. The stationary points of the system (3.24) and (3.25) satisfy the following conditions:

$$y=0 \qquad \text{and} \qquad \frac{\chi'(x)}{1+\chi(x)}=0, \quad (3.26)$$

i.e. $y=0$ and $\chi'(x)=0$ should be fulfilled if $\chi(x)\neq -1$. The stability conditions for the stationary point $(x,y)=(x_0,0)$ are given by

$$\left.\frac{\partial(dy/d\tau)}{\partial x}\right|_{(x,y)=(x_0,0)}=-\frac{\chi''(x_0)}{1+\chi(x_0)}<0, \quad (3.27)$$

$$\left.\frac{\partial(dy/d\tau)}{\partial y}\right|_{(x,y)=(x_0,0)}=-\sqrt{3\chi(x_0)}<0. \quad (3.28)$$

In case the scalar potential is positive, i.e. $\chi(x)>0$, the stability conditions (3.27) and (3.28) mean that $\chi''(x_0)>0$ is needed to provide stability.

As described above, there is not any points satisfying $dy/d\tau\to\pm\infty$ except for the cases $\chi\to\pm\infty$ or $\chi'\to\pm\infty$ when η is negative, whereas, there are nullclines defined by $dy/d\tau=0$.

Nullclines and attractors in the limit $x \to \pm\infty$

In general, the nullcline $dy/d\tau = 0$ is written as follows:

$$-\sqrt{3}y\left[1-y^2+\chi(x)\right]\sqrt{\left[\tfrac{1}{2}y^2+\chi(x)\right]\left(1-\tfrac{3}{2}y^2\right)}-\left(1-\frac{3}{2}y^2\right)\left(1-\frac{1}{2}y^2\right)\chi'(x)=0. \quad (3.29)$$

There is a special solution of Eq. (3.29),

$$y=\pm\sqrt{\frac{2}{3}}. \quad (3.30)$$

The other solutions of Eq. (3.29) are obtained by assuming the function form of $\chi(x)$. We will consider the case $\chi(\pm\infty)$ converges and the case $\chi(\pm\infty)$ goes to positive infinity. If $\chi(\pm\infty)$ converges, then $\chi'(\pm\infty) = 0$. Therefore, the solutions of Eq. (3.29) in the limit $x \to \pm\infty$ are only $y = 0$ and $y = \pm\sqrt{2/3}$. The stable line of them for the trajectories is only $y = 0$, so $y = 0$ is only the attractor in the limit $x \to \pm\infty$ if $\chi(\pm\infty)$ converges. The stability of the lines are evaluated by the sign of $dy/d\tau$ in the region $0 < |y| < \sqrt{2/3}$. On the other hand, the nullclines in the limit $|\chi(x)| \gg 1$ fulfill Eq. (3.30) or

$$y\chi^{\frac{3}{2}}(x) \simeq -\sqrt{3(1-\frac{3}{2}y^2)}\left(1-\frac{1}{2}y^2\right)\chi'(x). \quad (3.31)$$

If $y \neq \pm\sqrt{2/3}$, the behavior of the nullclines (3.31) are determined by the large/small relation between $\chi^{\frac{3}{2}}$ and $|\chi'|$. If $\chi^{\frac{3}{2}} \gg |\chi'|$ in the limit $x \to \pm\infty$, Eq. (3.31) yields

$$y=-\frac{\chi'(x)}{\sqrt{3}\chi^{\frac{3}{2}}(x)}. \quad (3.32)$$

Equation (3.32) is a stable curve for the trajectories, because $dy/d\tau$ is negative for positive y and $dy/d\tau$ is positive for negative y. However, Eq. (3.32) is not an asymptote of the trajectories, because it is seen that the nullcline (3.32) only exist in the second quadrant and in the fourth quadrant by taking into account $\chi'(+\infty) > 0$ ($\chi'(-\infty) < 0$) caused from $\chi(+\infty) \to +\infty$ ($\chi(-\infty) \to +\infty$). The nullclines (3.30) are unstable in this case, therefore, there is no attractors in the limit $x \to \pm\infty$ if $0 \neq |\chi'| \ll \chi^{\frac{3}{2}}$. Else if $\chi^{\frac{3}{2}} \ll |\chi'|$ holds, the asymptotes of the nullcline (3.31) is given by

$$y=-\sqrt{\frac{2}{3}}+\frac{9}{2\sqrt{6}}\frac{\chi^3(x)}{\chi'^2(x)}. \quad (3.33)$$

The curve (3.33) is stable in y direction for the trajectories, moreover, the trajectories on the curve go toward $x \to -\infty$ if $\chi(-\infty) \to +\infty$, because the nullcline (3.33) not only exist in the

fourth quadrant but also exist in the third quadrant. Therefore, Eq. (3.33) is the asymptote of the trajectories if $\chi^{\frac{3}{2}} \ll |\chi'|$. If $\chi' \propto \chi^{\frac{3}{2}}$, there is one solution of y in Eq. (3.31). The signature of the solution is negative, therefore, it also becomes the asymptote in the third quadrant. All kinds of the attractors in the case $\eta < 0$ is summarized in Table III.

The case $\eta = 0$

The case of $\eta = 0$ is equivalent to the quintessence model. We will shortly review the dynamics in this case. By defining the dimensionless variables,

$$x = \sqrt{\pi G}\phi, \quad y = \sqrt{\pi} G\,\dot{\phi}, \quad \tau = \frac{t}{\sqrt{G}}, \quad \chi(x) = \pi G^2 V(x/\sqrt{\pi G}), \tag{3.34}$$

we can rewrite Eq. (2.6) as the following autonomous dynamical system:

$$\frac{dx}{d\tau} = y, \tag{3.35}$$

$$\frac{dy}{d\tau} = -2y\sqrt{3(y^2 + 2\chi(x))} - \chi'(x). \tag{3.36}$$

In this case, $\chi(x)$ should less than 1 to hold the effective theory, because $\chi(x)$ is defined as $\chi(x) = \pi G^2 V(x/\sqrt{\pi G})$. Note that it is possible to satisfy both of the conditions $G^2V(\phi) \ll 1$ and $|\chi(x)| = 8\pi|\eta|GV(x/\sqrt{8\pi G}) \gg 1$ in the case $\eta \neq 0$. The stationary points of the system (3.35) and (3.36) satisfy the following conditions:

$$y = 0 \qquad \text{and} \qquad -\chi'(x) = 0. \tag{3.37}$$

The stability conditions for the stationary point $(x, y) = (x_0, 0)$ are given by

$$\left.\frac{\partial(dy/d\tau)}{\partial x}\right|_{(x,y)=(x_0,0)} = -\chi''(x_0) < 0, \tag{3.38}$$

$$\left.\frac{\partial(dy/d\tau)}{\partial y}\right|_{(x,y)=(x_0,0)} = -2\sqrt{6\chi(x_0)} < 0. \tag{3.39}$$

If $\chi(x_0)$ is positive, Eqs. (3.37)-(3.39) mean the local minimum of the potential is stable and the local maximum of the potential is unstable.

Equation (3.36) shows that the nullcline should satisfy $y\chi' < 0$ or $y = \chi' = 0$. Then the condition $\chi > 0$ impose the nullcline to be in the second quadrant or fourth quadrant in the limit $x \to \pm\infty$. Therefore, the nullclines cannot be the attractor in the region $x \to \pm\infty$

except for the case they are the stationary points. While, the nullcline $dy/d\tau = 0$ satisfy the following condition if $\chi(x) > 0$:

$$y^2 = -\chi(x) + \sqrt{\chi^2(x) + \frac{\chi'^2(x)}{12}}. \tag{3.40}$$

Equation (3.40) shows that there are one negative solution and one positive solution with respect to y. Therefore, there is only one nullcline in the limit $x \to +\infty$ ($x \to -\infty$). While, Eq. (3.36) says that $\mathrm{sgn}(dy/d\tau) = -\mathrm{sgn}(y)$ is satisfied in the limit $y^2 \gg \chi, |\chi'|$, so the nullcline is the only candidate for the attractor in the region $x \to \pm\infty$ or $y \to \pm\infty$ except for the stationary points. Because of the reasons above, there are no attractors in the region $x \to \pm\infty$ or $y \to \pm\infty$ except for the stationary points

IV. SPECIFIC EXAMPLES

In this section we apply the general consideration given above for some specific potentials.

A. Power-law potential

Let us consider the power-law potential

$$V(\phi) = M^4 \left(\frac{\phi}{\phi_0}\right)^\alpha, \tag{4.1}$$

where M and ϕ_0 are positive constants with dimension of mass, and α is a dimensionless real number. Note that cosmological dynamics in the theory of gravity with the scalar field possessing a nonminimal kinetic coupling to gravity and the power-law potential has been investigated in Ref. [23].

The potential (4.1) has only one local minimum $V(0) = 0$ if $\alpha > 0$. The condition that the point $\phi = x = 0$ and $\dot{\phi} = y = 0$ represents a stationary point is $V'(0) = 0$, i.e. $\alpha > 1$. The stability of such the minimum of the potential is shown in the previous section. Namely, it is stable if $V''(0) > 0$ regardless of the value of η. Therefore, the point $(x, y) = (0, 0)$ is always stable if it is a stationary point. While, the conditions seen in the previous section; $\chi(+\infty)$ is finite, $\chi(+\infty)$ is infinite and $|\chi'(+\infty)| \ll \chi^{1/2}(+\infty)$, $\chi(+\infty)$ is infinite and $|\chi'(+\infty)| \sim \chi^{1/2}(+\infty)$, and $\chi(+\infty)$ is infinite and $\chi^{1/2}(+\infty) \ll |\chi'(+\infty)| \ll \chi^{3/2}(+\infty)$, are translated into $\alpha \leq 0$, $0 < \alpha < 2$, $\alpha = 2$, and $2 < \alpha$, respectively. Therefore, there is



Figure 1: A phase diagram in the case $\chi(x) = x^2$. Solid gray curves and a dashed gray curve represent $dy/d\tau = 0$ and $dy/d\tau = \pm\infty$, respectively. Examples of the trajectories which go to the attractors are shown as colored curves in the left panel, and the corresponding time evolutions of the Hubble parameter $H(t)$ are expressed in the right panel.

one attractor $(x, y) = (+\infty, 0)$ in the limit $x \to +\infty$ if $\alpha < 0$ and $\eta > 0$. In the case $\alpha = 0$ and $\eta > 0$, there is one asymptote $y = \sqrt{8\pi G\eta M^4 - 1}$ if $8\pi G\eta M^4 > 1$, else if $8\pi G\eta M^4 < 1$, there is not asymptotes in the region $x \gg 1$. The line $y = -\chi'/[\sqrt{3\chi}(1-\chi)] = 0$ is not stable and is not unstable for x direction because $\chi' \equiv 0$. The solution on the asymptote $y = \sqrt{8\pi G\eta M^4 - 1}$ is $\phi(t) = \sqrt{(8\pi G\eta M^4 - 1)/8\pi G\eta}\ t + const.$, on the other hand, the solution on the line $y = 0$ is given by $\phi(t) = const.$. In the case $0 < \alpha < 2$ and $\eta > 0$, there are two asymptotes in the limit $x \to +\infty$: $y = \chi'(x)/\sqrt{3\chi^3(x)}$ and $y = \sqrt{\chi(x)}$. On the asymptote $y = \chi'(x)/\sqrt{3\chi^3(x)}$, the time evolution of $\phi = x/\sqrt{8\pi G}$ is given by

$$\phi(t) = \left[\frac{\alpha(\alpha+4)}{16\pi G\eta}\sqrt{\frac{\phi_0^\alpha}{24\pi G M^4}}t + const.\right]^{\frac{2}{\alpha+4}}. \tag{4.2}$$

Therefore, the Hubble rate on the asymptote is expressed as $H(t) \propto t^{\alpha/(4+\alpha)}$ in the limit $\phi_0 t \to +\infty$. This means so-called "little rip" expansion, which is the expansion faster than de Sitter expansion, is realized on the asymptote. The solution on the asymptote $y = \sqrt{\chi(x)}$

is expressed as

$$\phi(t) = \left(\frac{2-\alpha}{2} \sqrt{\frac{M^4}{\phi_0^\alpha}} t + const. \right)^{\frac{2}{2-\alpha}}. \tag{4.3}$$

Equation (4.3) shows that ϕ is an increasing function of time, but the Hubble rate is given by $H = 1/\sqrt{3\eta} = const.$ as shown in the previous section. In the case $\alpha = 2$ and $\eta > 0$, there are two asymptotes $y = \chi'(x)/\sqrt{3\chi^3(x)}$ and $y = \beta\sqrt{\chi(x)}$, $\beta \neq 0, \pm 1$ in the limit $x \to +\infty$. The behaviors of the asymptotes are almost same as those in the case $0 < \alpha < 2$, but $\phi(t)$ on the asymptote $y = \chi'(x)/\sqrt{3\chi^3(x)}$ becomes an exponential function of time. The coefficient β in $y = \beta\sqrt{\chi(x)}$ is determined by solving the equation $d[z\sqrt{\chi(x)}]/dx = (dy/d\tau)/(dx/d\tau)|_{y=z\sqrt{\chi(x)}}$ with respect to z. It can be explicitly written down as

$$rz^2\sqrt{2+z^2} = 2\mathrm{sgn}(z)(1-z^2), \tag{4.4}$$

where $r = \chi'(+\infty)/\sqrt{\chi(+\infty)} = 2\sqrt{\eta}M^2/\phi_0$. In general, there is one solution in Eq. (4.4) for positive r, and the solution β is located in the range $0 < \beta < 1$. In the case $\alpha > 2$ and $\eta > 0$, there are two asymptotes $y = \chi'(x)/\sqrt{3\chi^3(x)}$ and $y = \beta\sqrt{\chi^3(x)}/\chi'(x)$, $\beta \neq 0, \pm 2\sqrt{2}$ in the limit $x \to +\infty$. β in this case is easily determined by taking into account the condition $df(x)/dx = (dy/d\tau)/(dx/d\tau)$ on $y = f(x) = \beta\sqrt{\chi^3(x)}/\chi'(x)$ as

$$\beta = \frac{4\sqrt{2}\alpha}{3\alpha+2}. \tag{4.5}$$

Therefore, β is constrained in the range $\sqrt{2} < \beta < 4\sqrt{2}/3$. The solution on the asymptote $y = \beta\sqrt{\chi^3(x)}/\chi'(x)$ is given by

$$\phi(t) = \left(const. - \frac{\alpha\beta}{2}\sqrt{8\pi G}M^2 t \right)^{-\frac{2}{\alpha}} \phi_0. \tag{4.6}$$

The coefficient of t in Eq. (4.6) is negative, so the constant in the parenthesis should be positive. At some time, inside of the parenthesis in Eq. (4.6) vanishes, and $\phi(t)$ goes to be singular. The Hubble rate also goes to infinity because it on this attractor is given by $H = \sqrt{2V_\phi/(9\alpha(\eta V)^{1/2})}$. This type of singularity is called "Big Rip".

Summary of the expansion types in the limit $x \to +\infty$ is the following: de Sitter expansion is realized for $0 \leq \alpha \leq 2$, the Little Rip expansion is realized for $\alpha \geq 0$, and the Big Rip expansion is appeared for $\alpha > 2$. The case $\alpha = 2$ is depicted in Fig. 1. The damping oscillation of $H(t)$, de Sitter expansion, and the Little Rip expansion are realized on the asymptotes of the trajectories.



Figure 2: A phase diagram in the case $\chi(x) = 8(x^2 - 0.5)^2$. Solid gray curves and a dashed gray curve represent $dy/d\tau = 0$ and $dy/d\tau = \pm\infty$, respectively. Examples of the trajectories which go to the attractors are shown as colored curves in the left panel, and the corresponding time evolutions of the Hubble parameter $H(t)$ are expressed in the right panel.

B. Higgs-like potential

Now, let us consider a Higgs-like potential taking extremal values at $\phi = 0$ and $\phi = \pm\phi_0$:

$$V(\phi) = \frac{\lambda}{4}(\phi^2 - \phi_0^2)^2, \tag{4.7}$$

where λ is a positive dimensionless constant and ϕ_0 is a positive constant with dimension of mass. The detailed consideration of cosmological dynamics with the nonminimal kinetic coupling and the Higgs-like potential has been given in Ref. [24].

The stationary points located in the restricted region are given by $(x, y) = (0, 0), (\pm x_0, 0)$, where $x_0 = \sqrt{8\pi G}\phi_0$. Values of the potential at the stationary points are $V(\pm\phi_0) = 0$ and $V(0) = \lambda\phi_0^4/4$, i.e. $\chi(0) = 2\pi G|\eta|\lambda\phi_0^4$ if $\eta \neq 0$. The stationary points $(x, y) = (\pm x_0, 0)$ are stable, because the conditions $\chi(x_0) = 0$ and $\chi''(x_0) > 0$ are satisfied. On the other hand, the stability of the point $(x, y) = (0, 0)$ depends on the value of $\chi(0)$ in case $\eta > 0$. If $\chi(0) > 1$, then the point $(x, y) = (0, 0)$ is stable, otherwise it is unstable. A phase diagram in the case of $\chi(0) > 0$ is shown in Fig. 2. Note that the phase diagram is centrally symmetrical. In other words, the first (second) quadrant and the third (fourth) quadrant are completely

same except for the direction of the arrows are opposite, because the potential (4.7) is an even function. A behavior of trajectories in the limit $x \to \pm\infty$ is same as those in the case $\chi(x) \propto x^4$. There are two types of attractors in $x \to \pm\infty$; one of them causes the Little Rip expansion of the Universe, and another causes the Big Rip expansion. The numbers of attractors are seven in the whole phase diagram if $\chi(0) > 1$. However, the trajectories on the phase diagram are not so complicated, because the curve $dy/d\tau = \pm\infty$ divides the whole phase diagram into four pieces, and there are no trajectories that cross over the curve $dy/d\tau = \pm\infty$.

C. Exponential potential

In this section we consider the exponential potential

$$V(\phi) = M^4 \mathrm{e}^{-\phi/\phi_0}, \tag{4.8}$$

where M and ϕ_0 are positive constants with dimension of mass. If ϕ is not singular, then $V'(\phi)$ does not vanish and the exponential potential has no extremums. That is, there are no stationary points provided x is finite. The point $(x, y) = (+\infty, 0)$ is a stationary one. However, its stability analysis is quite difficult, because the relations $V(+\infty) = V'(+\infty) = V''(+\infty) = \cdots = 0$ are satisfied. Instead, we will use the analysis we have done in the region $|x| \gg 1$. Equation (4.8) yields $0 < \chi(x) \ll 1$ and $\chi'(x) < 0$ for $1 \ll |x| < +\infty$, so it is seen that the point $(x, y) = (+\infty, 0)$ is an attractor. To investigate a behavior of trajectories in the region $1 \ll x < +\infty$, we will impose the conditions $|y|, \chi, |\chi'| \ll 1$ on Eq. (3.3). Then we have

$$\frac{dy}{d\tau} \simeq -\sqrt{3}y\sqrt{\frac{1}{2}y^2 + \chi(x)} - \chi'(x). \tag{4.9}$$

It is seen that $dy/d\tau > 0$ is always satisfied if $y < 0$ by taking into account $\chi'(x) < 0$. Therefore, there is a stable nullcline in $y > 0$. This nullcline is the asymptote leading to the attractor $(x, y) = (+\infty, 0)$. The nullcline is given by

$$y = \sqrt{-\chi + \sqrt{\chi^2 + \frac{3}{2}\chi'^2}}. \tag{4.10}$$



Figure 3: A phase diagram in the case $\chi(x) = \mathrm{e}^{-x}$. Solid gray curves and a dashed gray curve represent $dy/d\tau = 0$ and $dy/d\tau = \pm\infty$, respectively. Examples of the trajectories which go to the attractors are shown as colored curves in the left panel, and the corresponding time evolutions of the Hubble parameter $H(t)$ are expressed in the right panel.

Taking into account that $y = dx/d\tau$ and $V(\phi) = M^4\mathrm{e}^{-\phi/\phi_0}$, we can resolve Eq. (4.10) with respect to $x(\tau)$ as follows

$$x(\tau) = 2x_0 \ln\left[\frac{1}{2x_0}\sqrt{\left(\sqrt{1+\frac{3}{2x_0^2}}-1\right)8\pi G\eta M^4\tau} + const.\right], \tag{4.11}$$

where $x_0 = \sqrt{8\pi G}\phi_0$. At the same time, the Hubble rate function is given as

$$H(t) = \frac{\gamma}{t}, \qquad \gamma^2 = \frac{2x_0^2\left(1+\sqrt{1+\frac{3}{2x_0^2}}\right)}{3\left(\sqrt{1+\frac{3}{2x_0^2}}-1\right)}. \tag{4.12}$$

If $\gamma = 1/2$, the Hubble rate expressed by Eq. (4.12) is same as that of the radiation dominant era.

On the other hand, there are the other attractors in the limit $x \to -\infty$. For the potential (4.8), $\chi(-\infty) \to +\infty$ and $|\chi'(-\infty)| \sim \chi(-\infty)$ are satisfied. Therefore, there are two kind of attractors for $x \to -\infty$ if $\eta > 0$; $y = \chi'/\sqrt{3\chi^3}$ and $y^2 \propto \sqrt{\chi^3}/\chi'$. Resolving

$x'(\tau) = \chi'(x(\tau))/\sqrt{3\chi^3(x(\tau))}$ with respect to $x(\tau)$ gives

$$x(\tau) = -2x_0 \ln\left[\frac{1}{4x_0}\sqrt{\frac{1}{6\pi G\eta M^4 x_0}}\tau + const.\right]. \tag{4.13}$$

Therefore, the Hubble rate parameter on the curve $y = \chi'/\sqrt{3\chi^3}$ is approximately given as

$$H(t) \simeq \frac{t}{6\eta x_0^{3/2}}. \tag{4.14}$$

Here, the constant in Eq. (4.13) is ignored. The time evolution given by Eq. (4.14) means that the Little Rip expansion is realized on the asymptote $y = \chi'/\sqrt{3\chi^3}$.

The coefficient of the curve $y^2 = \alpha\sqrt{\chi^3}/\chi'$ is obtained by equating $-d[\alpha\sqrt{\chi^3}/\chi']^{1/2}/dx$ with $(dy/d\tau)/(dx/d\tau)$ on the curve. Taking the conditions $|y| \sim \chi^{1/4} \sim |\chi'|^{1/4} \gg 1$ into account gives the following value of α:

$$\alpha = -\frac{4\sqrt{2}}{3} > -2\sqrt{2}. \tag{4.15}$$

Therefore, the concrete form of the asymptote is

$$y = -\sqrt{-\frac{4\sqrt{2}}{3}\frac{\chi^{3/2}(x)}{\chi'(x)}} = -4\sqrt{\frac{\sqrt{\pi G\eta M^4}x_0}{3}}\mathrm{e}^{-\frac{x}{4x_0}}. \tag{4.16}$$

Then, we can obtain the time evolution of x and H as follows:

$$x(\tau) = 4x_0 \ln\left[const. - \sqrt{\frac{\sqrt{\pi G\eta M^4}}{3x_0}}\tau\right], \tag{4.17}$$

$$H(t) = \frac{1}{const. - t}. \tag{4.18}$$

Eq. (4.18) means that the scale factor $a(t)$ and the Hubble rate $H(t)$ go to infinity at some finite time, i.e. the Big Rip occurs. These characteristic behaviors of the Hubble rate function can be seen in Fig. 3.

V. CONCLUSIONS

We have considered a cosmological dynamics of the scalar field with nonminimal kinetic coupling and a potential of general form. First, we have shown that the field equations can be represented as an autonomous system by using appropriate dimensionless parameters. It is worth noticing that the field equations expressed in a dimensionless form contains

ultimately only one arbitrary function $\chi(x) = 8\pi G|\eta|V(x/\sqrt{8\pi G})$ instead of two arbitrary values, that is the potential $V(\phi)$ and the coupling parameter η.

Stationary points of the autonomous system are expressed as $(x, y) = (x_0, 0)$ provided $\chi'(x_0) = 0$ and $\chi(x_0) \neq 1$. A stability of stationary points is determined by the value of $\chi(x_0)$ and the sign of $\chi''(x_0)$. In case $0 \leq \chi(x_0) < 1$ a stationary point is stable if $\chi''(x_0)$ is positive. In case $1 < \chi(x_0)$ it is stable if $\chi''(x_0)$ is negative. The other attractors of the system are $(x, y) = (\pm\infty, 0)$, $(\pm\infty, O(1))$, $(\pm\infty, \pm\infty)$. A behavior of asymptotes which go towards these attractors has been clarified by investigating critical lines and given in Sec. III. All results are arranged in Tables II and III. By using the results of Sec. III, we have shown that interesting scenarios of time evolution of the Universe, e.g. de Sitter expansion, the Little Rip and the Big Rip, are realized on the asymptotes given in Sec. IV. Existence conditions for the asymptotes depend in general on the form of $\chi(x)$, or, ultimately, $V(\phi)$.

In Sec. IV, some specific forms of the scalar potential $V(\phi)$ have been considered and a behavior of phase trajectories has been evaluated. As the result, it has been shown that the variety of asymptotes depends essentially on a power of ϕ. In particular, the Hubble rate on the asymptotes becomes the steeper the higher a power of ϕ. In the case of exponential potential, it has been shown that there exist generally three attractors which correspond to the decelerated expansion, the Little Rip, and the Big Rip, respectively.

Acknowledgments

The work was supported by the Russian Government Program of Competitive Growth of Kazan Federal University. SVS was supported by the RSF grant 16-12-10401.

[1] A. G. Riess *et al.* [Supernova Search Team Collaboration], Astron. J. **116**, 1009 (1998) [astro-ph/9805201].

[2] S. Perlmutter *et al.* [Supernova Cosmology Project Collaboration], Astrophys. J. **517**, 565 (1999) [astro-ph/9812133].

[3] E. Komatsu *et al.* [WMAP Collaboration], Astrophys. J. Suppl. **192**, 18 (2011) [arXiv:1001.4538 [astro-ph.CO]].

[4] P. A. R. Ade *et al.* [Planck Collaboration], Astron. Astrophys. **571**, A16 (2014) [arXiv:1303.5076 [astro-ph.CO]].

[5] P. A. R. Ade *et al.* [Planck Collaboration], arXiv:1502.01589 [astro-ph.CO].

[6] W. J. Percival *et al.* [SDSS Collaboration], Mon. Not. Roy. Astron. Soc. **401**, 2148 (2010) [arXiv:0907.1660 [astro-ph.CO]].

[7] C. Blake *et al.*, Mon. Not. Roy. Astron. Soc. **418**, 1707 (2011) [arXiv:1108.2635 [astro-ph.CO]].

[8] F. Beutler *et al.*, Mon. Not. Roy. Astron. Soc. **416**, 3017 (2011) [arXiv:1106.3366 [astro-ph.CO]].

[9] A. J. Cuesta *et al.*, Mon. Not. Roy. Astron. Soc. **457**, no. 2, 1770 (2016) [arXiv:1509.06371 [astro-ph.CO]].

[10] T. Delubac *et al.* [BOSS Collaboration], Astron. Astrophys. **574**, A59 (2015) [arXiv:1404.1801 [astro-ph.CO]].

[11] S. V. Sushkov, Phys. Rev. D**80**, 103505 (2009).

[12] E. N. Saridakis, S. V. Sushkov, Phys. Rev. D**81**, 083510 (2010).

[13] S. V. Sushkov, Phys. Rev. D**85**, 123520 (2012).

[14] S. Sushkov and R. Korolev, Class. Quant. Grav. **29**, 085008 (2012).

[15] A. A. Starobinsky, S. V. Sushkov, M. S. Volkov, JCAP 1606 (2016) no.06, 007.

[16] S. F. Daniel and R. R. Caldwell, Class. Quant. Grav. **24**, 5573 (2007). C. Gao, JCAP **06**, 023 (2010); L.N. Granda, W. Cardona, JCAP **1007**, 021 (2010); L. N. Granda, Class. Quantum Grav. **28**, 025006 (2011); L. N. Granda, JCAP **1104**, 016 (2011); L.N. Granda, E. Torrente-Lujan, J. J. Fernandez-Melgarejo, Eur. Phys. J. **C71**, 1704 (2011); L.N. Granda, arXiv:1109.1371; L.N. Granda, Mod. Phys. Lett. A **27**, 1250018 (2012); H. Mohseni Sadjadi, Phys. Rev. D **83**, 107301 (2011); A. Banijamali, B. Fazlpour, Phys. Lett. **B703**, 366 (2011); G. Gubitosi, E.V. Linder, Phys. Lett. **B703**, 113 (2011).

[17] C. Germani and A. Kehagias, Phys. Rev. Lett. **105**, 011302 (2010).

[18] C. Germani and A. Kehagias, Phys. Rev. Lett. **106**, 161302 (2011).

[19] S. Tsujikawa, Phys. Rev. D **85**, 083518 (2012).

[20] J. B. Dent, S. Dutta, E. N. Saridakis, and J.-Q. Xia, JCAP **11** (2013) 058.

[21] B. Ratra and P. J. E. Peebles, Phys. Rev. D **37**, 3406 (1988); C. Wetterich, Nucl. Phys. B **302**, 668 (1988); A. R. Liddle and R. J. Scherrer, Phys. Rev. D **59**, 023509 (1999); I. Zlatev, L. M. Wang and P. J. Steinhardt, Phys. Rev. Lett. **82**, 896 (1999); Z. K. Guo, N. Ohta and

Y. Z. Zhang, Mod. Phys. Lett. A **22**, 883 (2007); S. Dutta, E. N. Saridakis and R. J. Scherrer, Phys. Rev. D **79**, 103005 (2009).

[22] G. W. Horndeski, Int. J. Theor. Phys. 10, 363-384 (1974).

[23] M.A. Skugoreva, S.V. Sushkov, A.V. Toporensky, Phys. Rev. D**88**, 083539 (2013).

[24] J. Matsumoto and S. V. Sushkov, JCAP **1511**, no. 11, 047 (2015) [arXiv:1510.03264 [gr-qc]].

[25] A. A. Starobinsky, Sov. Astron. Lett. **4**, 82 (1978).